

AD-A096 654

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/6 12/1

AN INHOMOGENEOUS QUASILINEAR HYPERBOLIC SYSTEM.(U)

NOV 80 C WANG

DAA629-80-C-0041

UNCLASSIFIED

MRC-TSR-2137

NL

1 OF 1

AS A

Subend

END
DATE
FILMED
4 -81
DTIC

LEVEL II ②

AD A 096654

MRC Technical Summary Report # 2137

AN INHOMOGENEOUS QUASILINEAR
HYPERBOLIC SYSTEM

Ching-hua Wang

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

DTIC
SELECTED
MAR 23 1981
E

November 1980

(Received September 23, 1980)

Approved for public release
Distribution unlimited

81 3 19 065

FILE COPY

Sponsored by

U.S. Army Research Office
P.O. Box 12211
Research Triangle Park
North Carolina 27709

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

AN INHOMOGENEOUS QUASILINEAR HYPERBOLIC SYSTEM

Ching-hua Wang

Technical Summary Report # 2137

November 1980

ABSTRACT

1. Author	
2. Title	
3. Subtitle	
4. Author's Address	
5. Author's Phone	
6. Author's Fax	
7. Author's E-mail	
8. Author's Other	
9. Author's Other	
10. Author's Other	
11. Author's Other	
12. Author's Other	
13. Author's Other	
14. Author's Other	
15. Author's Other	
16. Author's Other	
17. Author's Other	
18. Author's Other	
19. Author's Other	
20. Author's Other	
21. Author's Other	
22. Author's Other	
23. Author's Other	
24. Author's Other	
25. Author's Other	
26. Author's Other	
27. Author's Other	
28. Author's Other	
29. Author's Other	
30. Author's Other	
31. Author's Other	
32. Author's Other	
33. Author's Other	
34. Author's Other	
35. Author's Other	
36. Author's Other	
37. Author's Other	
38. Author's Other	
39. Author's Other	
40. Author's Other	
41. Author's Other	
42. Author's Other	
43. Author's Other	
44. Author's Other	
45. Author's Other	
46. Author's Other	
47. Author's Other	
48. Author's Other	
49. Author's Other	
50. Author's Other	
51. Author's Other	
52. Author's Other	
53. Author's Other	
54. Author's Other	
55. Author's Other	
56. Author's Other	
57. Author's Other	
58. Author's Other	
59. Author's Other	
60. Author's Other	
61. Author's Other	
62. Author's Other	
63. Author's Other	
64. Author's Other	
65. Author's Other	
66. Author's Other	
67. Author's Other	
68. Author's Other	
69. Author's Other	
70. Author's Other	
71. Author's Other	
72. Author's Other	
73. Author's Other	
74. Author's Other	
75. Author's Other	
76. Author's Other	
77. Author's Other	
78. Author's Other	
79. Author's Other	
80. Author's Other	
81. Author's Other	
82. Author's Other	
83. Author's Other	
84. Author's Other	
85. Author's Other	
86. Author's Other	
87. Author's Other	
88. Author's Other	
89. Author's Other	
90. Author's Other	
91. Author's Other	
92. Author's Other	
93. Author's Other	
94. Author's Other	
95. Author's Other	
96. Author's Other	
97. Author's Other	
98. Author's Other	
99. Author's Other	
100. Author's Other	

We consider quasilinear hyperbolic partial differential equations modeling ideal gas flow under various physical effects. When these effects are represented as Lipschitz continuous functions of the states, solutions to the initial value problem are shown to exist globally in time. Our analysis is based on the random choice method which generalizes the Glimm scheme for hyperbolic conservation laws.

When the effects are strongly dissipative then the flow decays exponentially to a constant state as time tends to infinity.

AMS (MOS) Subject Classifications: 35L60, 35L67, 35B40

Key Words: Quasilinear, Initial value problem, Riemann Invariants, Weak solution, Existence of theorem of global solution, Asymptotic behavior, Inhomogeneous.

Work Unit Number 1 (Applied Analysis)

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

SIGNIFICANCE AND EXPLANATION

We study the initial value problem for quasilinear hyperbolic system of the following form:

$$\frac{\partial u}{\partial t} + \frac{\partial (\frac{1}{v})}{\partial x} = U(u, v, x, t)$$
$$\frac{\partial v}{\partial t} - \frac{\partial u}{\partial x} = V(u, v, x, t) .$$

The system models ideal gas flow where U and V represent physical effects such as damping and external moving force. Suppose that the initial data $(u(x,0), v(x,0))$ have bounded total variation, and U and V are Lipschitz continuous functions of u and v . Then the initial value problem has an admissible weak solution. When U and V represent strong dissipative effects, then the solution decays to a constant state as t tends to infinity. Our analysis is based on the random choice method which generalizes the Glimm scheme for conservation laws.

Our assumptions are general enough to include several important physical effects. Moreover, the random choice method can be applied to more general quasilinear hyperbolic systems as an effective numerical scheme.

The responsibility for the wording and views expressed in this descriptive summary lies with M.C., and not with the author of this report.

AN INHOMOGENEOUS QUASILINEAR HYPERBOLIC SYSTEM

Ching-hua Wang

INTRODUCTION

We consider the initial value problem for the inhomogeneous hyperbolic system

$$\begin{cases} \frac{\partial u}{\partial t} + \frac{\partial (\frac{1}{v})}{\partial x} = U(u, v, x, t) \\ \frac{\partial v}{\partial t} - \frac{\partial u}{\partial x} = V(u, v, x, t) \end{cases} \quad (0.1)$$

$$(u, v)(x, 0) = (u_0(x), v_0(x)) \quad (0.2)$$

Here $u_0(x)$ and $v_0(x)$ are bounded functions with bounded total variation, $v_0(x)$ has a positive lower bound. It is assumed that there exists a constant $M > 0$ such that U and V do not depend on x for $|x| > M$. U and V are smooth functions of u, v, x and t .

By the Riemann invariants

$$r = u + \log v \quad s = u - \log v \quad (0.3)$$

the system (0.1) can be diagonalized as

$$\begin{cases} \frac{\partial r}{\partial t} - \frac{1}{v} \frac{\partial r}{\partial x} = R(r, s, x, t) \\ \frac{\partial s}{\partial t} + \frac{1}{v} \frac{\partial s}{\partial x} = S(r, s, x, t) \end{cases} \quad (0.4)$$

where $R(r, s, x, t) = U + \frac{V}{v}$, $S(r, s, x, t) = U - \frac{V}{v}$; $(U, V) = (U, V)(\frac{r+s}{2}, \exp(\frac{r-s}{2}), x, t)$

Throughout this paper we also assume that there exist two constants K_1, K_2 such that

$$|R(r_1, s_1, x, t) - R(r_2, s_2, x, t)| \leq K_1(|r_1 - r_2| + |s_1 - s_2|) \quad (0.5)$$

$$|S(r_1, s_1, x, t) - S(r_2, s_2, x, t)| \leq K_1(|r_1 - r_2| + |s_1 - s_2|)$$

$$\left| \frac{\partial R}{\partial x} \right| \leq K_2, \quad \left| \frac{\partial S}{\partial x} \right| \leq K_2 \quad (0.6)$$

When $(U, V) \equiv (0, 0)$, system (0.1) reduces to the following system

$$\begin{cases} \frac{\partial u}{\partial t} - \frac{\partial(\frac{1}{v})}{\partial x} = 0 \\ \frac{\partial v}{\partial t} - \frac{\partial u}{\partial x} = 0 \end{cases} \quad (0.7)$$

The initial boundary value problem of (0.7) has been solved by Nishida T. [6]. When U and V do not depend on x and t , the global solution of (0.1 and (0.2) has been constructed by Ying Lung-an and Wang Ching-hua [7]. For general inhomogeneous system

$$\frac{\partial u}{\partial t} + \frac{\partial f(u)}{\partial x} = g(x, t, u) \quad (0.8)$$

Ying Lung-an and Wang Ching-hua [8] have proved the existence of the global solution of initial value problem of system (0.8) for $g(x, t, u) = g(u) e^{-Kt}$. When g does not depend on t , Liu Tai-Ping [5] has constructed global solutions for system (0.8) and studied their asymptotic behavior.

In general (0.1) and (0.2) does not possess smooth solutions, and we look for weak solutions in $0 \leq t < T$, i.e., solutions satisfying

$$\iint_{0 \leq t < T} (u \phi_t + \frac{1}{v} \phi_x + U \phi) dx dt + \int_{-\infty}^{+\infty} u_0(x) \phi(x, 0) dx = 0 \quad (0.9)$$

$$\iint_{0 \leq t < T} (v \psi_t - u \psi_x + V \psi) dx dt + \int_{-\infty}^{+\infty} v_0(x) \psi(x, 0) dx = 0 \quad (0.10)$$

for any smooth functions ϕ and ψ with compact support in $0 \leq t < T$.

The purpose of this paper is to prove the existence of the weak solution for (0.1) and (0.2) and study their asymptotic behavior as the time variable t tends to infinity.

1. Difference Scheme

We now describe the difference scheme we use for solving (0.1) and (0.2), which is the generalization of one in [2] and [7]. Randomly choose an equidistributed sequence $a=\{a_i\}$ in $(-1,1)$ and mesh length $\Delta x=l$ $\Delta t=h$ satisfying the Courant-Friedrich-Lewy condition

$$\frac{l}{h} = \text{const} \geq \frac{1}{v}$$

for all v under consideration.

Let $(\tilde{u}, \tilde{v})(x, t, m+1, k, u_l, v_l, u_r, v_r)$ denote the solution of the system (0.7) at the point $((m+1)l, kh)$ with the following initial data

$$(u, v)(x, kh) = \begin{cases} (u_l, v_l) & x < (m+1)l \\ (u_r, v_r) & x > (m+1)l \end{cases} \quad (1.1)$$

We take

$$(u, v)(x, t, m+1, k, u_l, v_l, u_r, v_r) = (\tilde{u} + U(\tilde{u}, \tilde{v}, (m+1)l, kh)(t-kh),$$

$$\tilde{v} \exp \left(\frac{V(\tilde{u}, \tilde{v}, (m+1)l, kh)}{\tilde{v}} (t-kh) \right) \quad (1.2)$$

as the approximation solution of the system (0.1) with the initial data (1.1). In fact, using the Riemann invariants, we can rewrite (1.2) in the following form

$$(r, s) = (\tilde{r} + R(\tilde{r}, \tilde{s}, (m+1)l, kh)(t-kh), \tilde{s} + S(\tilde{r}, \tilde{s}, (m+1)l, kh)(t-kh)) \quad (1.3)$$

The construction of the difference solution $(u,v)(x,t)$ proceeds as follows:

For each mesh length ℓ , let $(u_{\ell,0}(x), v_{\ell,0}(x))$ be defined by the equation

$$(u_{\ell,0}, v_{\ell,0})(x) = \begin{cases} (u_0(-\infty), v_0(-\infty)), & x < -(2[\frac{1}{\ell}]^2 + 1)\ell \\ (u_0(m\ell), v_0(m\ell)), & \text{for } |x| \leq (2[\frac{1}{\ell}]^2 + 1)\ell \text{ and} \\ & (m-1)\ell < x < (m+1)\ell, m=\text{even} \\ (u_0(+\infty), v_0(+\infty)), & x > ((2[\frac{1}{\ell}]^2 + 1)\ell \end{cases} \quad (1.4)$$

Then we set

$$(u,v)(x,t) = (u,v)(x,t, m+1,0, (u_{\ell,0}, v_{\ell,0})(m\ell), (u_{\ell,0}, v_{\ell,0})(m+2)\ell) \\ \text{for } m\ell < x < (m+2)\ell, 0 < t < h, m = \text{even} \quad (1.5)$$

Inductively suppose that $(u,v)(x,t)$ has been defined for $0 \leq t < kh$, we set

$$(u,v)(x,t) = (u,v)(x,t, m+1,k, (u,v)((m+a_k)\ell, kh-0), (u,v)((m+2+a_k)\ell, kh-0)) \\ \text{for } m\ell < x < (m+2)\ell, kh \leq t < (k+1)h, m+k = \text{even} \quad (1.6)$$

It is obvious that $(u,v)(x,t)$ depends on the mesh length h and sequence $a = \{a_i\}$, but we omit the subscript.

2. Bounds for the Difference Equation

Because we use (u, v) defined in (1.2) as the building block to build the difference solution of (0.1) and (0.2), we have to do some estimates on it.

First we consider $(\tilde{u}, \tilde{v}) (x, t, (m+1), k, u_\ell, v_\ell, u_r, v_r)$ defined in Section 1. It is well known that there are three constant region in the half plane $t > kh$, in which (\tilde{u}, \tilde{v}) equals to (u_ℓ, v_ℓ) , (u_m, v_m) and (u_r, v_r) respectively. (u_m, v_m) is connected to (u_ℓ, v_ℓ) by a 1-wave on the left and (u_r, v_r) is connected to (u_m, v_m) by a 2-wave on the left. We define (r_ℓ, s_ℓ) , (r_m, s_m) and (r_r, s_r) by means of (0.3) and define

$$\Delta r = r_m - r_\ell \quad \Delta s = s_r - s_m \quad (2.1)$$

$$P((u_\ell, v_\ell), (u_r, v_r)) = -\min(0, \Delta r) - \min(0, \Delta s) \quad (2.2)$$

It has been proved [6] that

$$P((u_i, v_\ell), (u_j, v_j)) < P((u_i, v_i), (u_k, v_k)) + P((u_k, v_k), (u_j, v_j)) \quad (2.3)$$

where u_i, u_k, u_j are any constants, v_i, v_k, v_j are any positive constants

Let

$$\delta r = r_r - r_\ell, \quad \delta s = s_r - s_\ell \quad (2.4)$$

we have

$$\begin{aligned}\delta r_t &= u_r + U_r(t-kh) + \log(v_r \exp \frac{v_r}{v_r}(t-kh)) - (u_l + U_l(t-kh) + \log(v_l \exp \frac{v_l}{v_l}(t-kh))) \\ &= (r_r - r_l) + (R_r - R_l)(t-kh)\end{aligned}$$

$$\begin{aligned}\delta s_t &= u_r + U_r(t-kh) - \log(v_r \exp \frac{v_r}{v_r}(t-kh)) - (u_l + U_l(t-kh) - \log(v_l \exp \frac{v_l}{v_l}(t-kh))) \\ &= (s_r - s_l) + (S_r - S_l)(t-kh)\end{aligned}$$

where U_r, V_r, R_r, S_r evaluated at $(u_r, v_r, (m+1)l, kh)$ and U_l, V_l, R_l, S_l evaluated at $(u_l, v_l, (m+1)l, kh)$.

Therefore by (0.5) we have

$$\begin{aligned}\text{Lemma 2.1} \quad & \begin{cases} |\delta r_t - \delta r| \leq K_1(|\delta r| + |\delta s|)(t-nh) \\ |\delta s_t - \delta s| \leq K_1(|\delta r| + |\delta s|)(t-nh) \end{cases} \quad (2.5)\end{aligned}$$

In the same way as Lemma 3 in [7], we have

Lemma 2.2 If (u_r, v_r) can be connected to (u_l, v_l) by only one wave, u_r^*, u_l^* are any constants and v_r^*, v_l^* are any positive constants then

$$P((u_l^*, v_l^*), (u_r^*, v_r^*)) \leq P((u_l, v_l), (u_r, v_r)) + 3(|\delta r^* - \delta r| + |\delta s^* - \delta s|) \quad (2.6)$$

where $\delta r^* = r_r^* - r_l^* = (u_r^* + \log v_r^*) - (u_l^* + \log v_l^*)$, $\delta s^* = (u_r^* - \log v_r^*) - (u_l^* - \log v_l^*)$.

In order to estimate the variations of $r(x, t)$ and $s(x, t)$, we define some functionals.

For integers $k \geq 1$, we define

$$F(kh+0) = \sum_{m+k=\text{even}} P((u, v)((m+a_k)l, kh-0), (u, v)((m+2+a_k)l, kh-0)),$$

and for integer $k \geq 0$

$$F((k+1)h-0) = \sum_{m+k=\text{even}} \{P((u,v)((m-1+a_{k+1})l, (k+1)h-0), \\ (u,v)(ml-0, (k+1)h-0)) +$$

$$P((u,v)(ml-0, (k+1)h-0), (u,v)(ml+0, (k+1)h-0)) +$$

$$P((u,v)(ml+0, (k+1)h-0), (u,v)((m+1+a_{k+1})l, (k+1)h-0))\}$$

Lemma 2.3. For any given $T > 0$, $\sum_{-\infty}^{+\infty} u(\cdot, t)$ and $\sum_{-\infty}^{+\infty} v(\cdot, t)$ the total variation of the difference solution $(u(x, t), v(x, t))$, are bounded uniformly for the mesh length h , sequence a and $t < T$, $v(x, t)$ has positive lower bound uniformly for the same parameters provided $k < \frac{1}{2K_1}$.

Proof. Since $U(u, v, x, t)$, $V(u, v, x, t)$ do not depend on x for $|x| > M$ and (0.5) there exists a constant \bar{M} depending only on T and $u_0(\pm\infty)$, $v_0(\pm\infty)$ such that $|s(\pm\infty, t)| \leq \bar{M}$, $|r(\pm\infty, t)| \leq \bar{M}$ for $0 \leq t < T$, provided h is small enough.

It follows from that $U(u, v, x, t)$, $V(u, v, x, t)$ do not depend on x for $|x| > M$ again and (0.6) that

$$\sum_{m+k=\text{even}} P((u,v)(ml-0, (k+1)h-0), (u,v)(ml+0, (k+1)h-0)) \leq$$

$$\leq h \sum_{m+k=\text{even}} \{|R((r,s)((m+a_k)l, kh-0), (m-1)l, kh) - R((r,s)(m+a_k)l, kh-0),$$

$$(m+1)l, kh)| + |S((r,s)((m+a_k)l, k-0), (m-1)l, kh) - S((r,s)(m+a_k)l, kh-0),$$

$$(m+1)l, kh)|\} \leq 4K_2 Mh.$$

It is obvious that $\int_{-\infty}^{+\infty} r(\cdot, nh+0) + \int_{-\infty}^{+\infty} s(\cdot, nh+0) \leq 4F(nh+0) + 4\bar{M}$.
Then from Lemma 2.1, Lemma 2.2, we obtain

$$\begin{aligned} F((n+1)h-0) &\leq F(nh+0) + 6K_1 h \left(\int_{-\infty}^{+\infty} r(\cdot, nh+0) + \int_{-\infty}^{+\infty} s(\cdot, nh+0) \right) + 4K_2 Mh \\ &\leq F(nh+0) (1+24K_1 h) + (24K_1 \bar{M}h + 4K_2 Mh) \end{aligned}$$

Because $F((n+1)h+0) \leq F((n+1)h-0)$ we have

$$F((n+1)h+0) \leq F(nh+0) (1+24K_1 h) + 24K_1 \bar{M}h + 4K_2 Mh$$

Let $F(0) = \sum_{m=\text{even}} P((u_0, v_0)(m\ell), (u_0, v_0)((m+2)\ell))$, then we obtain

$$F(0) \leq \int_{-\infty}^{+\infty} r_0(\cdot) + \int_{-\infty}^{+\infty} s_0(\cdot), \text{ where } r_0(x) = u_0(x) + \log v_0(x),$$

$$s_0(x) = u_0(x) - \log v_0(x).$$

After simple computation we get

$$F((n+1)h+0) \leq (F(0) + \bar{M} + \frac{K_2 M}{6K_1}) e^{24K_1 T}$$

$$\leq \left(\int_{-\infty}^{+\infty} r_0(\cdot) + \int_{-\infty}^{+\infty} s_0(\cdot) + \bar{M} + \frac{K_2 M}{6K_1} \right) e^{24K_1 T} \text{ for } (n+1)h < T$$

and thus

$$\begin{aligned} \int_{-\infty}^{+\infty} r(\cdot, nh+0) + \int_{-\infty}^{+\infty} s(\cdot, nh+0) &\leq 4 \left(\int_{-\infty}^{+\infty} r_0(\cdot) + \int_{-\infty}^{+\infty} s_0(\cdot) + \bar{M} + \frac{K_2 M}{6K_1} \right) e^{24K_1 T} + \\ &\quad + 4\bar{M} \text{ for } nh < T \end{aligned}$$

It follows from that $u_0(x)$ and $r_0(x)$ are bounded functions with bounded variation and $v_0(x)$ has a positive bound, Lemma 2.1 and $h < \frac{1}{2K_1}$ that

$$\int_{-\infty}^{+\infty} r(\cdot, t) + \int_{-\infty}^{+\infty} s(\cdot, t) \leq Q \quad \text{for } t < T \quad (2.7)$$

$$\text{where } Q = 8 \left(\int_{-\infty}^{+\infty} r_0(\cdot) + \int_{-\infty}^{+\infty} s_0(\cdot) + \bar{M} + \frac{K_2 M}{6K_1} \right) e^{24K_1 T} + \frac{K_2}{K_1} M + 8\bar{M}.$$

Therefore

$$|r(x, t)| \leq 2\bar{M} + Q, \quad |s(x, t)| \leq 2\bar{M} + Q, \quad |u(x, t)| \leq 2\bar{M} + \frac{Q}{2} \quad \text{for } t < T \quad (2.8)$$

$$e^{-(2\bar{M} + \frac{Q}{2})} \leq v(x, t) \leq e^{2\bar{M} + \frac{Q}{2}} \quad \text{for } t < T \quad (2.9)$$

$$\int_{-\infty}^{+\infty} u(\cdot, t) \leq Q/2 \quad \text{for } t < T \quad (2.10)$$

$$\int_{-\infty}^{+\infty} v(\cdot, t) \leq \frac{1}{2} Q e^{2\bar{M} + \frac{Q}{2}} \quad \text{for } t < T \quad (2.11)$$

Q.E.D.

Lemma 2.4. For any given $X > 0$, there exists constant D depending only on $U, V, u_0(x), v_0(x), T$ and X such that

$$I(X) = \int_{-X}^X (|u(x, t_2) - u(x, t_1)| + |v(x, t_2) - v(x, t_1)|) dx \leq \\ \leq D((t_2 - t_1) + h), \quad 0 < t_1, t_2 < T \quad (2.12)$$

provided h is small enough.

Proof: For definiteness suppose $t_1 < t_2$ and there is a pair of integers n, k such that $nh \leq t_1 < (n+1)h < \dots < (n+k)h < t_2 \leq (n+k+1)h$. It is obvious that

$$I(X) \leq I_1(X) + I_2(X)$$

where

$$\begin{aligned} I_1(X) &= \sum_{i=0}^{k+1} \int_{-X}^X (|u(x, (n+i)h+0) - u(x, (n+i)h-0)| + \\ &\quad + |v(x, (n+i)h+0) - v(x, (n+i)h-0)|) dx, \\ I_2(X) &= \sum_{i=0}^k \int_{-X}^X (|u(x, (n+i+1)h-0) - u(x, (n+i)h+0)| + \\ &\quad + |v(x, (n+i+1)h-0) - v(x, (n+i)h+0)|) dx \\ &\quad + \int_{-X}^X (|u(x, t_1) - u(x, nh+0)| + |v(x, t_1) - v(x, nh+0)|) dx + \\ &\quad + \int_{-X}^X (|u(x, t_2) - u(x, (n+k)h+0)| + |v(x, t_2) - v(x, (n+k)h+0)|) dx. \end{aligned}$$

It is not difficult to show that $I_1(X) \leq 2\ell \left(\left\lceil \frac{t_2 - t_1}{h} \right\rceil + 3 \right)$

$$\left\{ \sup_{0 < t < T} \sup_{-\infty}^{+\infty} u(\cdot, t) + \sup_{0 < t < T} \sup_{-\infty}^{+\infty} v(\cdot, t) \right\}$$

For $x \in ((m-1)\ell, (m+1)\ell)$, $m+n+i+1 = \text{even}$. Using (1.2) and (2.9) we have

$$|u(x, (n+i+1)h-0) - u(x, (n+i)h+0)| \leq \frac{(m+1)l}{(m-1)l} V u(x, (n+i)h+0) + |U^*| h.$$

$$|y(x, (n+i+1)h-0) - v(x, (n+i)h+0)| \leq \frac{(m+1)l}{(m-1)l} V r(x, (n+i)h+0) + 2|V^*| h$$

provided h is small enough, where

$$(U^*, V^*) = ((U, V), (x, (n+i+1)h-0), ml, (n+i)h).$$

For $\int_{-X}^X (|u(x, t_1) - u(x, nh+0)| + |v(x, t) - v(x, nh+0)|) dx$ and $\int_{-X}^X (|u(x, t_2) - u(x, (n+k)h+0)| + |v(x, t_2) - v(x, (n+k)h+0)|) dx$ we have similar estimate.

Combining the above estimates, we obtain

$$I_2 \leq \left(2 \sup_{0 < t < T} \sup_{-\infty}^{+\infty} V u(\cdot, t) + \sup_{0 < t < T} \sup_{-\infty}^{+\infty} V r(\cdot, t) \right) +$$

$$+ 2 X h \left(\sup_{m+n+i+1=\text{even}} U^* + 2 \sup_{m+n+i+1=\text{even}} V^* \right) \left[\left[\frac{t_2 - t_1}{h} \right] + 4 \right]$$

Since (2.8) (2.9) (2.10) and (2.11), we obtain (2.12). Q.E.D.

3. Convergence of the Difference Solutions

Because $(u, r)(x, t)$ is not an exact solution of (0.1) in the strip $kh \leq t < (k+1)h, k = 0, 1, \dots, \frac{T}{h}$, so in order to prove the existence of the weak solution of (0.1) and (0.2) we first prove two lemmas.

Lemma 3.1. If $x' = \xi_1 t'$ is a 1-shock wave of (\tilde{u}, \tilde{v}) $(x, t, (m+1)h, k, (u_\ell, v_\ell), (u_r, v_r))$ in the domain $\{(x, t) : m\ell < x < (m+2)\ell, kh \leq t < (k+1)h, m+k = \text{even}\}$ then

$$|\xi_1[u] - [\frac{1}{v}]| \leq wt'(|u_m - u_\ell| + |v_m - v_\ell|) \quad (3.1)$$

$$|\xi_1[v] - [u]| \leq wt'(|u_m - u_\ell| + |v_m - v_\ell|) \quad (3.2)$$

provided h is small enough, where $x' = x - (m+1)\ell, t' = t - kh$, the constant w depends only on $U, V, u_0(x), r_0(x); (u_\ell, v_\ell) = (u, v)((m+a_k)\ell, kh-0), (u_r, v_r) = (u, v)((m+2+a_k)\ell, kh-0); (u_m, v_m)$ is the constant solution being connected to (u_ℓ, v_ℓ) by 1-shock wave on the left; $[]$ denotes the jump of the quantity in bracket across the 1-shock wave.

For $x' = \xi_2 t'$ is a 2-shock wave of (\tilde{u}, \tilde{v}) , we have same kind of result.

Proof: It follows from that (u_r, v_r) and (u_m, v_m) satisfy Rankine-Hugoniot condition and (0.5) (2.8) (2.11) that

$$|\xi_1[u] - [\frac{1}{v}]| \leq \left| \xi_1 (u_m - u) t' - \left(\left(\frac{1}{v_m e^{\frac{v_m t'}{v_\ell}}} - \frac{1}{v_m} \right) - \left(\frac{1}{v_\ell e^{\frac{v_\ell t'}{v_\ell}}} - \frac{1}{v_\ell} \right) \right) \right| \leq$$

$$\begin{aligned}
&\leq |\xi_1 (U_m - U_\ell) t'| + \left| \frac{1}{v_\ell} \int_0^{\frac{v_\ell t'}{v_\ell}} e^{-\eta} d\eta - \frac{1}{v_m} \int_0^{\frac{v_m t'}{v_m}} e^{-\eta} d\eta \right| \leq \\
&\leq |\xi_1 (U_m - U_\ell) t'| + |(v_\ell - v_m) \left(-\frac{1}{v_\ell^2} \int_0^{\frac{v_\ell t'}{v_\ell}} e^{-\eta} d\eta + \frac{t'}{v_\ell} e^{-\frac{v_\ell t'}{v_\ell}} \frac{\partial}{\partial v} \left(\frac{v}{v_\ell} \right) \right)| + \\
&+ \left| \frac{1}{v_m} \int_0^{\frac{\tilde{v} t'}{v_m}} e^{-\eta} d\eta \right| \leq w t' (|u_m - u_\ell| + |v_m - v_\ell|)
\end{aligned}$$

where $(U_\ell, v_\ell) = (U, V)(u_\ell, v_\ell, (m+1)\ell, kh)$, $(U_m, v_m) = (U, V)(u_m, v_m, (m+1)\ell, kh)$

$v_\star = V(u_\ell, v_\star, (m+1)\ell, kh)$, $\tilde{v} = V(u_\ell, v_m, (m+1)\ell, kh)$;

$v_\star = \theta u_m + (1-\theta)v_\ell$, $0 < \theta < 1$. The proof of (3.2) is similar. Q.E.D.

Lemma 3.2. If (\tilde{u}, \tilde{v}) is a 1-rarefaction wave in the subdomain

$\Omega_1 = \{(x, t); \xi_0' < \frac{x'}{t'} < \xi_1', 0 < t' < h\}$ of the domain $\Omega = \{(x, t)$,

$m\ell < x < (m+2)\ell, kh \leq t < (k+1)h, m+k=\text{even}\}$; then

$$\begin{aligned}
&\left| \int_{\Omega_1} (u \phi_t + \frac{1}{v} \phi_x + U) dx dt + \int_{\partial \Omega_1} u \phi ds - \frac{1}{v} \phi dt \right| \\
&\leq R_1 h^3 + R_2 h^2 \left(\frac{\xi_1' t'}{\xi_0' t'} u(kh+0) + \frac{\xi_1' t'}{\xi_0' t'} v(\cdot, kh+0) \right) \quad (3.3)
\end{aligned}$$

$$\begin{aligned}
&\left| \int_{\Omega_1} (v \chi_t - u \chi_x + V) dx dt + \int_{\partial \Omega_1} v \chi dx - u \chi dt \right| \\
&\leq R_1 h^3 + R_2 h^2 \left(\frac{\xi_1' t'}{\xi_0' t'} u(\cdot, kh+0) + \frac{\xi_1' t'}{\xi_0' t'} v(\cdot, kh+0) \right) \quad (3.4)
\end{aligned}$$

provided h is small enough.

Where $x' = x - (m+1)l$, $t = t - kh$; $(\tilde{u}, \tilde{v}) = (\tilde{u}, \tilde{v})(x, t, (m+1)l, k, (u, v)(m+1)l, kh-0), (u, v)((m+2+a_k)l, kh-0))$, ϕ and χ are smooth functions with compact support in $0 \leq t < T$; R_1 and R_2 are constants depending only on $U, V, u_0(x), r_0(x), \phi$ and χ , $(U, V) = (U(u, v, x, t), V(u, v, x, t))$.

For (\tilde{u}, \tilde{v}) is a 2-rarefaction wave in the subdomain $\Omega_2 = \{(x, t); \xi'_1 < \frac{x'}{t} < \xi_2, 0 < t < h\}$ of the domain Ω , we have same kind of result.

Proof: (\tilde{u}, \tilde{v}) , as a 1-rarefaction wave in Ω_1 , satisfies (0.7) in Ω_1 in classic sense. Because of (0.5) and (2.9), U and V , as the functions of u and v , satisfy Lifshitz condition with the Lifshitz constant depending only on K_1, \bar{M} and Q .

It follows from (1.2)(1.6) that

$$\frac{\partial u}{\partial x} = \frac{\partial \tilde{u}}{\partial x} + \frac{\partial \tilde{U}}{\partial \tilde{u}} \frac{\partial \tilde{u}}{\partial x} + \frac{\partial \tilde{U}}{\partial \tilde{v}} \frac{\partial \tilde{v}}{\partial x}) t' \quad (3.5)$$

$$\frac{\partial u}{\partial t} = \frac{\partial \tilde{u}}{\partial t} + \frac{\partial \tilde{U}}{\partial \tilde{u}} \frac{\partial u}{\partial t} + \frac{\partial \tilde{U}}{\partial \tilde{v}} \frac{\partial \tilde{v}}{\partial t}) t' + \tilde{U} \quad (3.6)$$

$$\begin{aligned} \frac{\partial v}{\partial x} = \frac{\partial \tilde{v}}{\partial x} e^{\frac{\tilde{V} t'}{\tilde{v}}} + (\tilde{v})^{-1} t' e^{\frac{\tilde{V} t'}{\tilde{v}}} (\tilde{v} \frac{\partial \tilde{v}}{\partial u} \frac{\partial \tilde{u}}{\partial x} + \\ + \tilde{v} \frac{\partial \tilde{V}}{\partial \tilde{v}} \frac{\partial \tilde{v}}{\partial x} - \tilde{V} \frac{\partial \tilde{v}}{\partial x}) \quad (3.7) \end{aligned}$$

$$\frac{\partial v}{\partial t} = \frac{\partial \tilde{v}}{\partial t} e^{\frac{(\tilde{v}t')}{\tilde{v}}} + (\tilde{v})^{-1} t' e^{\frac{(\tilde{v}t')}{\tilde{v}}} \left(\tilde{v} \frac{\partial \tilde{v}}{\partial \tilde{u}} \frac{\partial \tilde{u}}{\partial t} + \tilde{v} \frac{\partial \tilde{v}}{\partial \tilde{v}} \frac{\partial \tilde{v}}{\partial t} - \tilde{v} \frac{\partial \tilde{v}}{\partial t} \right) + \tilde{v} e^{\frac{(\tilde{v}t')}{\tilde{v}}} \quad (3.8)$$

where $(\tilde{U}, \tilde{V}) = (\tilde{U}, \tilde{V})(\tilde{u}, \tilde{v}, (m+1)l, kh)$.

Let $J_1(x, t) = \frac{\partial u}{\partial t} + \frac{\partial(\frac{1}{v})}{\partial x} - U$, then we obtain

$$\begin{aligned} J_1(x, t) &= \frac{\partial \tilde{u}}{\partial t} + \left(\frac{\partial \tilde{u}}{\partial \tilde{u}} \frac{\partial \tilde{u}}{\partial t} + \frac{\partial \tilde{u}}{\partial \tilde{v}} \frac{\partial \tilde{v}}{\partial t} \right) t' + \tilde{U} - (\tilde{v})^{-2} e^{-\frac{(\tilde{v}t')}{\tilde{v}}} \\ &\quad \left(\frac{\partial \tilde{v}}{\partial x} e^{\frac{(\tilde{v}t')}{\tilde{v}}} + (\tilde{v})^{-1} t' e^{\frac{(\tilde{v}t')}{\tilde{v}}} \left(\tilde{v} \frac{\partial \tilde{v}}{\partial \tilde{u}} \frac{\partial \tilde{u}}{\partial x} + \tilde{v} \frac{\partial \tilde{v}}{\partial \tilde{v}} \frac{\partial \tilde{v}}{\partial x} - \tilde{v} \frac{\partial \tilde{v}}{\partial x} \right) - \right. \\ &\quad \left. U = (1 - e^{\frac{(\tilde{v}t')}{\tilde{v}}}) (\tilde{v})^{-2} \frac{\partial \tilde{v}}{\partial x} + t' \left(\frac{\partial \tilde{v}}{\partial \tilde{u}} \frac{\partial \tilde{u}}{\partial t} + \frac{\partial \tilde{u}}{\partial \tilde{v}} \frac{\partial \tilde{v}}{\partial t} \right) - t' (\tilde{v})^{-3} e^{\frac{(\tilde{v}t')}{\tilde{v}}} \right. \\ &\quad \left. \left(\tilde{v} \frac{\partial \tilde{v}}{\partial \tilde{u}} \frac{\partial \tilde{u}}{\partial x} + \tilde{v} \frac{\partial \tilde{v}}{\partial \tilde{v}} \frac{\partial \tilde{v}}{\partial x} - \tilde{v} \frac{\partial \tilde{v}}{\partial x} \right) \right). \end{aligned}$$

Using (0.5) and (0.6) for estimating $\tilde{U} - U$, then we get

$$\begin{aligned} &\iint_{\Omega_1} (u \phi_t + \frac{1}{v} \phi_x + U \phi) dx dt + \int_{\partial \Omega_1} u \phi dx - \frac{1}{v} \phi dt = \iint_{\Omega} J_1(x, t) \phi(x, t) dx dt \leq \\ &\leq R_1 h^3 + R_2 h^2 \left(\frac{\xi_1 t'}{v} u(\cdot, kh+0) + \frac{\xi_1 t'}{v} v(\cdot, kh+0) \right). \end{aligned}$$

The proof of (3.4) is similar.

Q.E.D.

We use the difference scheme introduced in section 2 and take the ratio of the mesh lengths $\frac{l_i}{h_i} = e^{2\bar{M}+Q}$ and $h_i = \frac{T}{2^i}$. When i is large enough, the difference solution $(u, v)(x, t)$ can be defined in $0 \leq t < T$.

The difference solution $(u, v(x, t))$ is discontinuous at the segment $\{(x, t): x = m\ell_i, kh_i < t \leq (k+1)h_i, m+k=\text{even}\}$. But it follows from (0.6), (2.8) and (2.11) that

$$\left| \int_{kh_i}^{(k+1)h_i} (u(m\ell_i+0, t) - u(m\ell_i-0, t)) \chi dt \right| \leq \int_{kh_i}^{(k+1)h_i} (t - kh_i) |U((u, v)((m+a_k)\ell_i, (m+1)\ell_i, kh_i) - U((u, v)((m+a_k)\ell_i, (m-1)\ell_i, kh_i)) \chi dt| \leq E h_i^3 \quad (3.9)$$

$$\left| \int_{kh_i}^{(k+1)h_i} \left(\frac{1}{v(m\ell_i+0, t)} - \frac{1}{v(m\ell_i-0, t)} \right) \phi dt \right| \leq E h_i^3 \quad (3.10)$$

provided i is large enough, where E is a constant depending only on K_2 , \bar{M} and Q .

With Lemma 3.1, lemma 3.2 and (3.9), (3.10) in mind by the way similar to these in the section 4 in [7] we obtain the following existence theorem.

Theorem 3.1. Suppose that $u_0(x)$ are bounded functions with bounded variation, $v_0(x)$ has a positive lower bound. $U(u, v, x, t)$, $V(u, v, x, t)$ are smooth functions satisfying (0.5) (0.6) and do not depend on x as $|x| > M$, where M is any given positive number, then for any given $T > 0$ the initial value problem (0.1) and (0.2) has a weak solution in $0 < t < T$.

4. Asymptotic behavior

Now we consider the asymptotic behavior of the solution of (0.1) and (0.2) under the following additional restrictions on U and V :

$$R \text{ depends only on } r, \quad \text{and} \quad R'(r) \leq -\alpha; \quad (4.1)$$

$$S \text{ depends only on } s, \quad \text{and} \quad S'(s) \leq -\alpha \quad (4.2)$$

where α is a positive number.

Lemma 4.1. Suppose that $R(r)$ satisfies (4.1), then the initial value problem

$$\begin{cases} \frac{dr}{dt} = R(r) \\ r(0) = r_0 \end{cases} \quad (4.3)$$

$$r(0) = r_0 \quad (4.4)$$

possesses a unique solution $r(t, r_0)$ satisfying

$$|r(t, r_0) - \bar{r}| \leq |r_0 - \bar{r}| e^{-\alpha t} \quad \text{for } t > 0 \quad (4.5)$$

where \bar{r} is the unique solution of $R(r) = 0$; and for

$$r_h(t, r_0) = \begin{cases} r_0 + R(r_0)t & , \quad 0 \leq t \leq h \\ r_h(h, r_0) + R(r_h(h, r_0))(t-h) & , \quad h \leq t \leq 2h \\ \dots & \\ r_h(nh, r_0) + R(r_h(nh, r_0))(t-nh) & , \quad nh \leq t \leq (n+1)h \\ \dots & \end{cases} \quad , \quad \text{the Euler's}$$

difference solutions of (4.3) (4.4), the following inequality holds

$$|r_h(t, r_0) - \bar{r}| \leq |\bar{r} - r_0| e^{-\alpha t} \quad \text{for } t > 0 \quad (4.6)$$

provided $h < \frac{1}{\alpha}$.

Proof. Since the assumptions (0.5), (4.3) and (4.4) possesses a unique solution $r(t, r_0)$ for all $t > 0$. By (4.1) we have

$$\frac{d(r(t, r_0) - \bar{r})}{dt} = R(r(t, r_0)) = -\alpha(r(t, r_0) - \bar{r})$$

and therefore (4.5).

Now we turn to prove (4.6).

Without loss of generality we assume $r_0 \neq \bar{r}$, because of $r_h(t, \bar{r}) \equiv r(t, \bar{r})$.

For definiteness suppose $r_0 > \bar{r}$. It follows from (0.5) (4.1) that for any $\tilde{r} > \bar{r}$ and $t_0 > 0$ the initial value problem

$$\begin{cases} \frac{dr}{dt} = R(r) \\ r|_{t=t_0} = \tilde{r} \end{cases} \quad \text{has a unique solution } r(t, \tilde{r}, t_0) \text{ such that}$$

$$\frac{d^2 r(t, \tilde{r}, t_0)}{dt^2} > 0; \text{ therefore}$$

$$\tilde{r} < r_h(t, r_0) < r(t, r_0) \quad \text{for } t > 0 \quad (4.7)$$

provided $h < \frac{1}{\alpha}$.

Similary for $r_0 < \bar{r}$ we have

$$r(t, r_0) < r_h(t, r_0) < \bar{r} \quad \text{for } t > 0 \quad (4.8)$$

By (4.5) (4.7) (4.8) we obtain (4.6).

Lemma 4.2. Under the assumption (4.1) and (4.2) we have

$$F(k+1)h-0) \leq (1-(1-c)ah)F(kh+0), \text{ for } k \geq 0 \quad (4.9)$$

provided $h < \min(\frac{1}{\alpha} \frac{1}{2K_1})$, where c is a constant depending only on $\lim_{x \rightarrow -\infty} u_0(\cdot)$, $\lim_{x \rightarrow -\infty} v_0(\cdot)$, the positive lower bound of $v_0(x)$ and the equation (0.7).

Proof: Now we consider $(\tilde{u}, \tilde{v})(x, t, (m+1), k, u_\ell, v_\ell, u_r, v_r)$ defined in section 1 again and first remind some results in [2] and [7].

There is a c^2 function $f(x)$ satisfying

$$\begin{aligned} f(x) &= \text{constant} & \text{for } x \geq 0 \\ 0 < f'(x) < 1, f''(x) < 0, & \text{for } x < 0 \\ \lim_{x \rightarrow -\infty} f'(x) &= 1 \end{aligned} \quad (4.10)$$

such that (u_m, v_m) is connected to (u, v) by 1-wave if and only if

$$s_m - s_\ell = f(r_m - r_\ell) \quad (4.11)$$

where $r_m - r_\ell \geq 0$ means that the 1-wave is a 1-rarefaction wave, and $r_m - r_\ell < 0$ means that the 1-wave is a 1-shock wave; and (u_r, v_r) is connected to (u_ℓ, v_ℓ) by 2-wave on the left if and only if

$$v_r - r_m = f(s_r - s_m) \quad (4.12)$$

$s_r - s_m \geq 0$ ($s_r - s_m < 0$) means that the 2-wave is a 2-rarefaction wave (a 2-shock wave respectively).

By the notation of (2.1) and (2.4) we have

$$\delta r = \Delta r + f(\Delta s) , \quad \delta s = \Delta s + f(\Delta r) \quad (4.13)$$

Because of (4.10) we get

$$\Delta r = p(\delta r, \delta s) , \quad \Delta s = q(\delta r, \delta s) \quad (4.14)$$

where $p, q \in C^2$. It is easy to know from simple computation that

$$\left\{ \begin{array}{ll} \frac{\partial p}{\partial(\delta r)} = \frac{1}{1-f'(p)f'(q)} & \frac{\partial q}{\partial(\delta r)} = \frac{-f'(p)}{1-f'(p)f'(q)} \\ \frac{\partial p}{\partial(\delta s)} = \frac{-f'(q)}{1-f'(p)f'(q)} & \frac{\partial q}{\partial(\delta s)} = \frac{1}{1-f'(p)f'(q)} \end{array} \right. \quad (4.15)$$

Suppose that (u_m, v_m) is connected to (u_ℓ, v_ℓ) by a 1-shock wave on the left. Since (0.5)(1.3) and (4.1), then we have

$$\left\{ \begin{array}{lll} s_m - s_\ell < (s_m + hS(s_m)) & -(s_\ell + hS(s_\ell)) & < 0 \\ r_m - r_\ell < (r_m + hR(r_m)) & -(r_\ell + hR(r_\ell)) & < 0 \end{array} \right. ,$$

$$\text{therefore } P((u_{\ell,h}, v_{\ell,h}), (u_{m,h}, v_{m,h})) < P(u_\ell, v_\ell), (u_m, v_m) \quad (4.16)$$

$$\text{where } u_{i,h} = u_i + U(u_i, v_i)h \quad v_{i,h} = v_i \exp\left(\frac{V(u_i, v_i)h}{v_i}\right) ,$$

i can be ℓ , m , or r .

Suppose that (u_m, v_m) is connected to (u_ℓ, v_ℓ) by a 1-rarefaction wave on the left. Since (0.5)(1.3) and (4.1), then we have

$$s_m + hS(s_m) = s_\ell + S(s_\ell)$$

$$0 < r_m + hR(r_m) - (r_\ell + hR(r_\ell))$$

therefore

$$P((u_{\ell,h}, v_{\ell,h}), (u_{m,h}, v_{m,h})) = P((u_\ell, v_\ell), (u_m, v_m)) = 0 \quad (4.17)$$

The same kinds of results as (4.16) and (4.17) can be obtained for a 2-shock wave and a 2-rarefaction wave in (\tilde{u}, \tilde{v}) respectively.

By combining the estimates (4.16), (4.17) and (2.3) we have

$$F((k+1)h-0) \leq F(kh+0) \quad \text{for } k \geq 0 \quad (4.18)$$

therefore for $\lim_{V \rightarrow +\infty} r(\cdot, t)$ and $\lim_{V \rightarrow +\infty} s(\cdot, t)$, the decreasing variation of $r(x, t)$ and $s(x, t)$ on the interval $(-\infty, +\infty)$, the following inequalities hold

$$\begin{aligned} \lim_{V \rightarrow +\infty} r(\cdot, t) &\leq F(0) \leq \left(\lim_{V \rightarrow +\infty} r_0(\cdot) + \lim_{V \rightarrow +\infty} s_0(\cdot) \right) = Q_1 \\ \lim_{V \rightarrow +\infty} s(\cdot, t) &\leq F(0) \leq \left(\lim_{V \rightarrow +\infty} r_0(\cdot) + \lim_{V \rightarrow +\infty} s_0(\cdot) \right) = Q_1 \end{aligned} \quad (4.19)$$

It is not difficult to know from (4.20) that the strength of the shock wave in (\bar{u}, \bar{v}) is less than Q_1 , i.e.

$$|\Delta r| \leq Q_1 \quad \text{for} \quad 1\text{-shock wave} \quad (4.20)$$

$$|\Delta s| \leq Q_1 \quad \text{for} \quad 2\text{-shock wave} \quad (4.21)$$

Since (4.10) then

$$0 \leq c = f'(-Q_1) < 1 \quad (4.22)$$

Now we are ready to obtain the refinement of (4.16) suppose that (u_m, v_m) is connected to (u_ℓ, v_ℓ) by a 1-shock wave on the left. By (4.15) we have

$$\begin{aligned} P((u_{\ell,h}, v_{\ell,h}), (u_{m,h}, v_{m,h})) - P((u_\ell, v_\ell), (u_m, v_m)) = \\ \int_{\delta r}^{\delta r_h} \frac{1-f'(p)}{1-f'(p)f'(q)} d\xi + \int_{\delta s}^{\delta s_h} \frac{1-f'(q)}{1-f'(p)f'(q)} d\eta \geq (1-c)((\delta r_h - \delta r) + \\ + (\delta s_h - \delta s)) \geq (1-c)h [(R(r_m) - R(r_\ell) + (S(s_m) - S(s_\ell)))] \\ \geq - (1-c)\alpha h \Delta r \end{aligned} \quad (4.23)$$

where $\delta r_k = (r_m + hR(r_m)) - (r_\ell + hR(r_\ell))$

$$\delta s_h = (s_m + hS(s_m)) - (s_\ell + hS(s_\ell))$$

Suppose that (u_r, v_r) is connected to (u_m, v_m) by a 2-shock wave on the left. Similarly we have

$$\begin{aligned}
 P((u_{m,h}, v_{m,h}), (u_{r,h}, v_{r,h})) - P((u_m, v_m), (u_r, v_r)) &\geq \\
 &\geq -(1-c)\alpha h \Delta s \quad (4.24)
 \end{aligned}$$

By combining the estimates (4.23) (4.24) (4.17) and (2.3), we have

$$F(k+1)h-0 \leq (1-(1-c)\alpha h)F(kh+0) \quad . \quad \text{Q.E.D.}$$

We turn now to proving the existence of the weak solution of (0.1) and (0.2) and studying its asymptotic behavior. We use $(u,v)(x,t)$ to denote the difference solutions of (0.1) and (0.2) before for convenience, although the difference solutions of (0.1) and (0.2) depend on the mesh length h and the sequence a . In order to distinguish between the difference solutions of (0.1) and (0.2) and the weak solution of (0.1) and (0.2), from now on we use $(u^h, v^h)(x,t)$ to denote the difference solutions of (0.1) and (0.2) and $(u,v)(x,t)$ to denote the weak solution of (0.1) and (0.2).

Theorem 4.1. Suppose that $u_0(x), v_0(x)$ are bounded functions with bounded variation and do not depend x as $|x| > M$, $v_0(x)$ has a positive lower bound. U, V are smooth function satisfying (0.5), (0.6) (4.1) and (4.2) then the weak solution $(u,v)(x,t)$ of (0.1) and (0.2) exists in $t > 0$ and satisfies the following decay laws.

$$\lim_{t \rightarrow \infty} u(\cdot, t) \leq 2Q_1 e^{-(1-c)at} + Q_2 e^{-at} \quad (4.25)$$

$$\lim_{t \rightarrow \infty} \log v(\cdot, t) \leq 2Q_1 e^{-(1-c)at} + Q_2 e^{-at} \quad (4.26)$$

$$\left| u(x, t) - \frac{\bar{r} + \bar{s}}{2} \right| \leq 2Q_1 e^{-(1-c)at} + \frac{3}{2} Q_2 e^{-at} \quad (4.27)$$

$$\left| \log v(x, t) - \frac{\bar{r} - \bar{s}}{2} \right| \leq 2Q_1 e^{-(1-c)at} + \frac{3}{2} Q_2 e^{-at} \quad (4.28)$$

where $Q_1 = \lim_{t \rightarrow \infty} r_0(\cdot) + \lim_{t \rightarrow \infty} s_0(\cdot)$, $Q_2 = \frac{1}{2}(|r_0(-\infty) - \bar{r}| + |r_0(+\infty) - \bar{r}| + |s_0(-\infty) - \bar{s}| + |s_0(+\infty) - \bar{s}|)$, \bar{r} and \bar{s} are the unique solution of $R(r) = 0$ and $S(s) = 0$ respectively, $c = f'(-Q_1)$.

Proof. If we choose the ratio of the mesh lengths

$$\frac{l}{h} = \text{const} \geq e^{\frac{r^* + s^*}{2}} + 2Q_1$$

where $r^* = \max(r_0(-\infty), \bar{r})$, $s^* = \max(s_0(-\infty), \bar{s})$, and $h = 2^{-i}$, where i is large enough such that $h < \min(\frac{1}{\alpha}, \frac{1}{2K_1})$, then the difference solution $(u^h, v^h)(x, t)$ defined in section 1 can be defined in all half plan $t > 0$ because that $|r^h(-\infty, t)| \leq r^*$, $|s^h(-\infty, t)| \leq s^*$ by Lemma 4.1 and $\lim_{t \rightarrow \infty} r^h(\cdot, t) \leq 2Q_1$, $\lim_{t \rightarrow \infty} s^h(\cdot, t) \leq 2Q_1$ by (4.19). By Lemma 3.1 and Lemma 3.2 we can obtain the existence of the weak solution $(u, v)(x, t)$ of (0.1) and (0.2) in $t > 0$ in similar way to that in theorem 3.1.

Using (4.9) and $F(kh+0) \leq F(kh-0)$, $k = 1, 2, \dots$, we obtain

$$F(nh+0) \leq (1-(1-c)ah)^n F(0) = (1-(1-c)ah)^n Q_1,$$

then thus

$$\begin{cases} \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} r^h(\cdot, nh+0) \leq (1-(1-c)ah)^n Q_1 \\ \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} s^h(\cdot, nh+0) \leq (1-(1-c)ah)^n Q_1 \end{cases} \quad (4.29)$$

It is not difficult by the means similar to this in Lemma 4.2 to know that

$$\begin{cases} \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} r^h(\cdot, t) \leq \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} r^h(\cdot, nh+0) \\ \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} s^h(\cdot, t) \leq \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} s^h(\cdot, nh+0) \end{cases}, \quad \text{for } nh < t < (n+1)h. \quad (4.30)$$

let i tends to infinite, by (4.29) we have

$$\begin{cases} \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} r(\cdot, t+0) \leq Q_1 e^{-(1-c)at} \\ \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} s(\cdot, t+0) \leq Q_1 e^{-(1-c)at} \end{cases} \quad \begin{matrix} \text{for } t = j_2^{-k} \text{ and } k, j \text{ are any} \\ \text{positive integers} \end{matrix} \quad (4.31)$$

Therefore by (4.30) and (4.31) we obtain

$$\begin{cases} \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} r(\cdot, t) \leq Q_1 e^{-(1-c)at} \\ \begin{matrix} +\infty \\ DV \\ -\infty \end{matrix} s(\cdot, t) \leq Q_1 e^{-(1-c)at} \end{cases} \quad \begin{matrix} \text{for } t > 0 \end{matrix} \quad (4.32)$$

It follows from (4.6) that

$$|r^h(-\infty, t) - \bar{r}| \leq |r_0(-\infty) - \bar{r}| e^{-at}, \quad |r^h(+\infty, t) - \bar{r}| \leq |r_0(+\infty) - \bar{r}| e^{-at}$$

$$|s^h(-\infty, t) - \bar{s}| \leq |s_0(-\infty) - \bar{s}| e^{-at}, \quad |s^h(+\infty, t) - \bar{s}| \leq |s_0(+\infty) - \bar{s}| e^{-at}$$

therefore

$$|r(-\infty, t) - \bar{r}| \leq |r_0(-\infty) - \bar{r}| e^{-at}, \quad |r(+\infty, t) - \bar{r}| \leq |r_0(+\infty) - \bar{r}| e^{-at}$$

$$|s(-\infty, t) - \bar{s}| \leq |s_0(-\infty) - \bar{s}| e^{-at}, \quad |s(+\infty, t) - \bar{s}| \leq |s_0(+\infty) - \bar{s}| e^{-at}.$$

Thus then

$$\left\{ \begin{array}{l} \int_{-\infty}^{+\infty} r(\cdot, t) \leq 2Q_1 e^{-(1-c)at} + (|r_0(-\infty) - \bar{r}| + |r_0(+\infty) - \bar{r}|) e^{-at} \\ \int_{-\infty}^{+\infty} s(\cdot, t) \leq 2Q_1 e^{-(1-c)at} + (|s_0(-\infty) - \bar{s}| + |s_0(+\infty) - \bar{s}|) e^{-at} \\ |r(x, t) - \bar{r}| \leq \int_{-\infty}^{+\infty} r(\cdot, t) + |r_0(-\infty) - \bar{r}| e^{-at} \\ |s(x, t) - \bar{s}| \leq \int_{-\infty}^{+\infty} s(\cdot, t) + |s_0(-\infty) - \bar{s}| e^{-at} \end{array} \right. \quad (4.33)$$

The estimates (4.25), (4.26), (4.27) and (4.28) follow from (4.33).

Q.E.D.

5. Entropy Condition

The entropy condition for quasilinear hyperbolic systems of conservation laws has been extensively studied by Lax, P.D. [3], Dafermos, C.M [1] and Liu, Tai-Ping [4]. Here we will prove a theorem about entropy condition for the weak solution of (0.1) and (0.2) obtained in Section 3.

Theorem 5.1. Under the same assumptions of theorem 3.1, the weak solution $(u,v)(x,t)$ of (0.1) and (0.2) constructed in theorem 3.1 satisfies the following entropy condition:

$$\iint_{0 < t < T} (g_t E + g_x F + g(uU - \frac{1}{v} V)) dx dt \geq 0 \quad (5.1)$$

for any non-negative smooth function g with compact support in $0 < t < T$; where $E(u,v) = \frac{u^2}{2} - \log v$, $F(u,v) = \frac{u}{v}$.

Proof. Suppose (u_m, v_m) is connected to (u_l, v_l) by a 1-shock wave with the speed of propagation s_1 on the left in the solution of (0.7), let $G(v_m, (u_l, v_l)) = s_1(E(u_m, v_m) - E(u_l, v_l)) - (F(u_m, v_m) - F(u_l, v_l))$, then we have

$$G(v_l, (u_l, v_l)) = 0$$

$$\frac{\partial}{\partial v_m} G(v_m, (u_l, v_l)) > 0$$

because $s_1 < 0$, $v_m > v_l$ for a 1-shock wave in the weak solution of (0.7).

Therefore

$$G(v_m, (u_l, v_l)) > 0 \quad (5.2)$$

Suppose (u_r, v_r) is connected to (u_m, v_m) by a 2-shock wave with the speed of propagation s_2 on the left in the solution of (0.7). Let $G(v_r, (u_m, v_m)) = s_2(E(u_r, v_r) - E(u_m, v_m)) - (F(u_r, v_r) - F(u_m, v_m))$, then similarly we have

$$G(v_r, (u_m, v_m)) > 0 \quad (5.2')$$

Using (3.3) (3.4) (3.9) and (3.10), we have

$$\iint_{t < 0 < T} (g_t E^h + g_x F^h + g(u^h u^h - \frac{v^h}{v^h}) dx dt = O(h) + I_1 + I_2 \quad (5.3)$$

Here

$$I_1 = \sum_{n=1}^{(\frac{T}{h})} g(x, nh) (E(u^h, v^h)(x, nh-0) - E(u^h, v^h)(x, nh+0)) dx \quad (5.4)$$

$$I_2 = \sum_{\substack{m+n=\text{even} \\ i=1,2}} \int_0^h g(s_{m,n,i} [E^h] - [F^h]) dt' \quad (5.5)$$

where $s_{m,n,i}$ is the speed of propagation of the i -shock wave in $(\tilde{u}, \tilde{v})(x, t(m+1), n, (u^h, v^h)(m+a_n)l, nh-0), (u^h, v^h)((m+2+a_n)l, nh-0)$, $t' = t - nh$. $[\cdot]$ denotes the jump of the quantity in the bracket across the shock wave. $E^h = E((u^h, v^h)(x, t))$, $F^h = F((u^h, v^h)(x, t))$.

Using (3.1) (3.2), we have

$$I_2 = O(h) + I'_2 \quad (5.6)$$

$$\text{where } I'_2 = \sum_{\substack{m+n=\text{even} \\ i=1,2}} \int_0^h g(s_{m,n,i} (E(\tilde{u}, \tilde{v})) - (F(\tilde{u}, \tilde{v}))) dt'$$

By (5.1), (5.2), we have

$$I'_2 \geq 0 \quad (5.7)$$

The entropy condition (5.1) follows from (5.3) (5.4) (5.5) (5.6) and (5.7).

Q.E.D.

References

1. Dafermos, C.M.: The entropy rate admissibility criterion for solutions of hyperbolic conservation laws. *J. Diff. Eq.* 14, 202-212 (1973).
2. Glimm, J.: Solutions in the large for nonlinear hyperbolic systems of equations. *Comm. Pure Appl. Math.* 27, 377-405 (1974).
3. Lax, P.D.: Shock waves and entropy. *Proc. Symp. Univ. Wisc.*, 1971, E.H. Zarantonell, ed., 603-634.
4. Liu, T.-P.: The entropy condition and the admissibility of shock. *Math. Anal. and Appl.* 53, 78-88 (1976).
5. Liu, T.-P.: Quasilinear hyperbolic systems. *Commun. Math. Phys.* 68, 141-172 (1979).
6. Nishida, T.: Global solutions for an initial bounded value problem of a quasilinear hyperbolic system. *Proc. Japan Acad.* 44, 462-466 (1968).
7. Ying Lung-an and Wang Ching-hua: Global solutions of the Cauchy problem for a nonhomogeneous quasilinear hyperbolic system.
8. Ying Lung-an and Wang Ching-hua: Solutions in the large for nonhomogeneous quasilinear hyperbolic systems of equations.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER MR2-TSR-2137	2. GOVT ACCESSION NO. AD-A096	3. RECIPIENT'S CATALOG NUMBER 654
4. TITLE (and Subtitle) An Inhomogeneous Quasilinear Hyperbolic System		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) Ching-hua/Wang		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1 (Applied Analysis)
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Technical summary Repts.		12. REPORT DATE Nov 1980
		13. NUMBER OF PAGES 31
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Quasilinear, Initial value problem, Riemann Invariants, Weak solution, Existence of theorem of global solution, Asymptotic behavior, Inhomogeneous.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We consider quasilinear hyperbolic partial differential equations modeling ideal gas flow under various physical effects. When these effects are represented as Lipschitz continuous functions of the states, solutions to the initial value problem are shown to exist globally in time. Our analysis is based on the random choice method which generalizes the Glimm scheme for hyperbolic conservation laws.		

20. ABSTRACT, continued,

When the effects are strongly dissipative then the flow decays exponentially to a constant state as time tends to infinity.

DATE
FILMED
-8